

D Departement U W Umweltwissenschaften



Swimming competition! Who beats the flow in a hydraulic barrier for invasive fish?

<u>Joschka Wiegleb¹</u>, Philipp E. Hirsch¹, Frank Seidel², Georg Rauter³, Patricia Burkhardt-Holm¹

Round goby is an invasive species, which outcompetes native species and represents a major threat for invaded ecosystems. Fish passes are bottlenecks that must be passed by every fish migrating upstream. The proposed hydraulic barrier is designed to create hydraulic conditions that impede passage of the round goby,

The prototype hydraulic barrier

- Stainless steel plate (3 mm) of 1m length to extend the jet stream and exceed the swimming capacity of the round goby
- ✓ Diagonal side walls to affect the flow predominantly at the bottom
- ✓ Smooth surface to prevent the round goby from substrate

whilst allowing passage of native species.

To test the barrier, ca. 43 fish of every species were released at the water outlet of a vertical slot fish pass model. The fish were free to swim upstream for two hours, unaffected by human presence. The species were tested separately.

Gudgeon experienced significantly smaller hydraulic forces on the barrier than the other



Prototype hydraulic barrier Flow

Methods

- > Measurement of hydraulic forces experienced by preserved fish
- Recording of swimming trajectories using cameras
- Comparing the trajectories betweem species using Artificial Intelligence

Trajectory

Round goby



(Neogobius melanostomus Pallas, 1814) is native in the Black and Caspian Sea. Round goby arrived in the river Rhine near Basel (Switzerland) in 2012, outcompete native species, and continue dispersal upstream.

Gudgeon



(Gobio gobio L.) is a poweful, semi-pelagic swimmer with similar ecological niche as round goby and bullhead.

holding

Figure1: **Hydraulic forces** experienced by 7 preserved fish per species over the barrier at 14 measurement points.



Bullhead



(Cottus gobio L.) looks similar to the round goby, inhabits small tributaries, and has a burst -and-hold swimming style.

Bullh

80L/s105 L/s130 L/s Predominantly round No round goby but six Round goby and goby but no gudgeon gudgeon passed the barrier. gudgeon passed the passed the barrier. Three **bullhead** passed and barrier. immediately returned.

Artificial Intelligence

Random Forest models revealed distinct swimming behaviour between species as they were able to predict the species of round goby and gudgeon on the swimming trajectories with high F1 scores.

Gudgeon Round goby

85.48 0.00 22.22

True





Figure. 2: The **swimming trajectories** of the three species on the barrier were extracted from the video footage of all three tested water discharges by hand. The 130 L/s water discharge is representative for real fish passes. The gray lines represent barrier contours.

Figure. 3: Crossvalidation of the Random Forest model, which was trained to predict the species on the swimming trajectories of Fig. 2. The confusion matrix displays prediction proportions [%] for the different species with correct prediction in the gray areas. A perfect model has an F1 score of 1.

Wiegleb, J., Hirsch, P. E., Seidel, F., Rauter, G., & Burkhardt-Holm, P. (2022). Flow, force, behaviour: assessment of a prototype hydraulic barrier for invasive fish. *Hydrobiologia*, 849(4), 1001–1019. https://doi.org/10.1007/s10750-021-04762-z



¹Program Man-Society-Environment, Department of Environmental Science, University of Basel, Vesalgasse 1, 4051 Basel, Switzerland ²Institute of Water and River Basin Management, Karlsruhe Institute of Technology, Kaiserstrasse 12, 76131 Karlsruhe, Germany ³BIROMED-Lab, Department ob Biomedical Engineering, University of Basel, Gewerbestrasse 14, 4123 Allschwil, Switzerland

Round

goby